

## WARNINGS AND PRECAUTIONS

- Keep out of the reach of children, uninformed persons and animals.
- May cause slight eye irritation.
- May cause allergic reactions in sensitive persons.
- Do not use after expiry date.
- Reseal packaging after use.
- Do not mix in containers that have previously been used for toxic substances.
- Wash mixing containers well with soap and water before use.
- Shake or stir the product mixture frequently during use to avoid settling out of the product.
- Avoid inhaling the dust.
- Wash hands with soap and water before eating, drinking or smoking.
- Do not use under extreme conditions i.e. temperatures above 35°C or at soil/water pH less than 4 or higher than 7.

See enclosed directions for application and use



CAUTION  
VERSIGTIG



# Eco-T<sup>®</sup>

(L6938)

Reg. No. L6938, Act No. 36 of 1947 / N-AR1047

GROUP

BM02

FUNGICIDE

A fungal inoculant for the suppression of crop root diseases and for enhanced growth of a wide variety of vegetables, crops, ornamentals and Eucalyptus species.



 **Andermatt**  
Madumbi



250g, 1kg, 5kg

## Active ingredient

*Trichoderma asperellum* strain kd (Min 2 x 10<sup>9</sup> spores per gram)



Founding Member:



Inspected by ECOCERT SA F-32600

Product suitable for use in organic agriculture complying to the annexes of the (EC) regulation n° 834/2007 and 889/2008 and NOP Regulation

Registered and produced by:

**Andermatt PHP (Pty) Ltd.**

Co. Reg. No. 2003/007987/07

Country of origin: South Africa



Distributed in South Africa by:

**Andermatt Madumbi (Pty) Ltd**

Co. Reg. No. 2009/020798/07

Tel: +27 (0) 33 342 3984 | support@andermatt.co.za

Suite 105, 24 Hilton Avenue, Hilton, 3245

[www.andermatt.co.za](http://www.andermatt.co.za)



15/06/2021



Batch No and Expiry date:

PEEL

# Eco-T®

Reg. No. L6938, Act No. 36 of 1947 / N-AR1047

GROUP

BM02

FUNGICIDE

**A fungal inoculant for the suppression of crop root diseases and for enhanced growth of a wide variety of vegetables, crops, ornamentals and Eucalyptus species.**

**Active ingredient:** *Trichoderma asperellum* strain kd  
(Min 2 x 10<sup>9</sup> spores per gram)

**DISCLAIMER:** Although this product has been tested under a variety of conditions, the supplier does not warrant that it will be efficacious under all conditions. Because the active agent is a living organism, its action may be affected by storage, handling, climatic or soil conditions. The supplier is also not responsible for poor performance of the product due to failure of the user to follow the label instructions. This naturally-occurring organism should not cause damage to crops, vegetation or the environment, nor harm to man and animals. However, in the unlikely event of this happening, the supplier does not accept responsibility for any damage that may occur.

## IMPORTANT POINTS TO REMEMBER

- Remember you are working with a living organism.
- Store at 4 – 10°C for maximum shelf-life.
- Do not leave the powder or mixture standing in the sun.
- Use within 6 hours of mixing.
- When using irrigation systems a minimum dilution of 1 kg/100 L is recommended.

## COMPATIBILITY

- Do not mix with any agrochemical without first consulting your sales agent.

## DIRECTIONS FOR USE - USE ONLY AS DIRECTED.

### SEED TREATMENT

- Seed dressing application rates differ for different crops.
- Apply to maize at a rate of 1 g/kg of seed or 25-50 g/ha; to wheat at 1-2 g/kg seed or 40-80 g/ha; or to soybeans at 1 g/kg seed or 50-75 g/ha.
- For rates on other crops please consult your local agent or Madumbi Sustainable Agriculture - [www.madumbi.co.za](http://www.madumbi.co.za) for details.
- A sticker is not required where *Eco-T*® is added during on-farm treatment with insecticides, compatible fungicides, or Rhizobium inoculant.
- Apply as a dry powder to slightly moist seed.

## DRENCH

- Apply as a drench to seedling trays, seedbeds, potted plants and turf at planting, and monthly thereafter.
- Mix at a rate of 1g/4L water.
- 250 g of this product treats 3000 m<sup>2</sup> of seedbed or 3000 seedling trays.
- Use 250-500 g/ha for field applications.
- Mix the required amount of the powder to a thin paste with a small amount of water before adding the remaining water.
- Irrigate the product into the root zone after application.