

WARNINGS AND PRECAUTIONS

- Keep out of reach of children, uninformed persons and animals.
- Avoid contact with skin and eyes.
- Do not eat, drink or smoke in areas where product is used.
- Wash hands and other exposed areas with soap and water before leaving work.
Remove contaminated clothing and shoes. Wash clothing and equipment after handling.
- May irritate skin and eyes. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.
- Clean applicator before and after use. Do not contaminate other crops, grazing, rivers, dams and areas not under treatment with wash water.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsing water to the contents of the spray or mixing tank.
Destroy the container, and do not use empty container for any other purpose.

See enclosed directions for application and use.



 **Andermatt**
Madumbi



200g



MycoUp Activ[®]

FERTILISER GROUP III

Reg. No. M0010, Act No. 36 of 1947

A biological fertiliser containing active mycorrhizal fungi to support healthy root development.

BIOSTIMULANT



Active ingredient

Glomus iranicum var. *tenuihypharum* var. *nova*.
(min. 1.2×10^4 spores per ml)

Manufactured by:
Symborg SL.
CIF: B73574832



Symborg

Founding member:



Registered and distributed by: **Andermatt Madumbi (Pty) Ltd**
Co. Reg. No. 2009/020798/07
Country of origin: South Africa
Tel: +27 (0) 33 342 3984
support@andermatt.co.za
Suite 105, 24 Hilton Avenue, Hilton, 3245
www.andermatt.co.za

DRAFT

15.06.2021



Batch No:
Expiry date:

PEEL

MycoUp Activ®

Reg. No. M0010, Act No. 36 of 1947

A biological fertiliser containing active mycorrhizal fungi to support healthy root development.

Active ingredient: *Glomus iranicum* var. *tenuihypharum* var. *nova*. (min. 1.2×10^4 spores per ml)

DISCLAIMER: Although this product has been tested under a variety of conditions, the supplier does not warrant that it will be efficacious under all conditions. Because the active agent is a living organism, its action may be affected by storage, handling, climatic or soil conditions. The supplier is also not responsible for poor performance of the product due to failure of the user to follow the label instructions. This naturally-occurring organism should not cause damage to crops, vegetation or the environment, nor harm to man and animals. However, in the unlikely event of this happening, the supplier does not accept responsibility for any damage that may occur.

STORAGE AND HANDLING

- Store in original container in a dry area, away from food and feed stuff.
- *MycoUp Activ®* can be stored at room temperature for up to 24 months.
- Use before expiry date indicated on label.
- Wear protective clothing, rubber gloves, goggles and respirator when handling.
- Do not leave the product or mixture standing in the sun.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED.

MIXING INSTRUCTIONS

- Measure required amount of *MycoUp Activ®* and dissolve in a small volume of water.
- Fill the application tank with half the required volume of water then add the *MycoUp Activ®* premix and mix well.
- If one is present, start agitator mechanism. Maintain agitation during application.
- Top up spray tank and check final pH is 5-8.
- If other products are to be tank mixed always add them first and correct pH.

IMPORTANT POINTS TO REMEMBER

- Apply as a soil drench application using conventional irrigation equipment, good coverage of the root zone is essential.
- It may be necessary to follow application with additional irrigation to ensure product penetrates into the soil profile.
- Mixed volumes must be applied as soon as possible and not left overnight or standing in the sun.
- Avoid applying *MycoUp Activ®* during peak sunlight hours.

COMPATIBILITY

- Compatibility with chemical products is not assured. Do not mix with chemicals unless the supplier has confirmed compatibility. Do not mix in containers that have previously been used for toxic substances.

Crop	Dosage	Remarks
Horticultural crops (greenhouse, field, hydroponics)	2-3 kg/ha	<ul style="list-style-type: none">• Apply 7 days after transplant.• For crops with an extended growing cycle repeat application at 2 kg/Ha 70-90 days after first application.
Strawberry	2-3 kg/ha	<ul style="list-style-type: none">• Apply 20 days after transplant.
Woody crops (citrus, pome and stone fruit, tree nuts, grapevines etc.)	2-3 kg/ha	<ul style="list-style-type: none">• Apply at early flowering, adjust rate accordingly to age of crop with higher rate on fully productive orchards.• Repeat application may be done annually.

NOTE: *MycoUp Activ®* should NOT be applied to the following crops *Chenopodiaceae* (e.g. Spinach), *Cruciferaeae* (cole crops such as cabbage, cauliflower, broccoli), *Fumariaceae*, *Cyperaceae*, *Commelinaceae*, *Urticaceae* (nettles) and *Poligonaceae* (buckwheat, rhubarb).