



elephant vert

fytoSave



Preventative biofungicide for stimulation of defenses on crops as specified



COMPOSITION:

COS-06A (Oligosaccharide complex) : 12,5g/l
Fungicide Group : P4 (Resistance Not Known)
Registration number : L10052

FORMULATION TYPE:

Soluble concentrate (SL)

DIRECTIONS FOR USE. USE ONLY AS DIRECTED

RESISTANCE MANAGEMENT:

Currently there is no known resistance to **FytoSave®** or the biological active. Due to the nature of the mode of action resistance is not expected to be likely. It is still advisable to use good agricultural practices of resistance management.

WAITING PERIOD AND USE RESTRICTION :

FytoSave® does not impose any risk of residues on harvested crops.

TO DELAY FUNGICIDE RESISTANCE:

- Avoid exclusive repeated use of fungicides from the same fungicide group code.
- Alternate or tank mix with products from different fungicide group codes.
- Refer to individual product labels when alternating products or when using tank mixtures with products in fungicide group codes 7, 27, 28, 29, 30 and M.
- **FytoSave®** may be applied alone or in combination and/ or rotation with other control methods (chemical, cultural, biological) as a tool for integrated disease management.

WARNINGS AND PRECAUTIONS:

- Avoid direct contact with eyes and skin.
- In case of accidental contact wash affected area with clean water.
- Wear gloves when handling.
- Wash hands before consuming food, drinking or smoking.
- Do not inhale spray mist. Do not eat, drink or smoke whilst mixing or applying.
- Prevent contamination of food, foodstuffs, drinking water and eating utensils.
- Clean applicator before and after use.
- Store in original container in a dry area, away from food and feed stuffs.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsing water to the contents of the spray or mixing tank. Destroy the container, and do not use empty container for any other purpose.
- Keep out of reach of children, uninformed persons and animals.
- Store the product in a cool, dry place.
- Shake container well before use.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label as well as by the methods, time and accuracy of application. The registration holder, furthermore, does not accept responsibility for damage to crops vegetation, the environment or harm to man or animals, the lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of registration. Please consult the supplier in the event of any uncertainty.

REGISTERED USAGES:

| CROPS / PEST | DOSAGES | REMARKS |
|--|-------------------------------------|---|
| Cucurbits Powdery mildew (<i>Podosphaera fusca</i>) | 3 L/ha (water volume 500-1000 L) | <ul style="list-style-type: none"> • Apply when disease is first noticed or as a precautionary application when conditions favour disease outbreak. • Repeat application every 7 days until disease is under control or conditions are no longer favourable. • Apply as a full cover film spray using conventional spray equipment. Good coverage, preferably of the lower leaf, is essential. |
| Table Grapes Downy mildew (<i>Plasmopara viticola</i>) Powdery mildew (<i>Erysiphe necator</i>) | 3 L/ha (water volume 200-1000 L) | <ul style="list-style-type: none"> • Apply as a full cover film spray using conventional spray equipment. Good coverage, preferably of the lower leaf, is essential. |

MIXING INSTRUCTIONS:

- Fill spray tank with half the required water volume and check neutral pH. Use an appropriate buffering agent if necessary.
- Start agitator mechanism. Maintain agitation during application.
- Add the correct amount of product to the water in the tank at the recommended dose rate.
- Top up spray tank and check final pH.
- If other products are to be tank mixed always add them first.
- Mixed volumes must be applied as soon as possible and not left overnight or standing in the sun.

REGISTERED BY

FytoFend (Pty) Ltd
Block A, Inanda Greens Business Park,
98 Albertyn Avenue, Sandton, 2196
Tel: +27 14 940 0083
Email: info@fytofend.com
Co Reg. 2018/306062/07

PRODUCED BY:

FytoFend S.A
Rue Georges Legrand, 6
B5032, Isnes, Belgium

PROUDLY DISTRIBUTED BY:

Andermatt Madumbi (Pty) Ltd
Co. Reg. No. 2009/020798/07
Tel: +27 (0) 33 342 3984
Email: support@andermatt.co.za
24 Hilton Ave, Suite 105,
Hilton, 3245



CAUTION
VERSIKTIG

